cold of high latitudes. I have found the flowers a second time formed, and shooting from the ground in November. The strong and unpleasant odour which every part of the plant emits on being broken, and which is precisely similar to that of the *Viverra mephitis;* has given it by an almost common consent, in every part of the country, the appellations of *Skunk weed* and *Skunk cabbage.*

The structure of this singular vegetable has caused it successively to be assigned to the genera *Arum*, *Dracontium* and *Pothos*, with none of which it fully agrees. Of the Aroideæ, to which it is related, it approaches most nearly in its flower to *Pothos;* while its fruit has more affinity to *Orontium.* The Rev. Dr. Cutler many years ago, in the Transactions of the American Academy, pointed out the distinctive characters of this plant, and pronounced it a new genus. No name, however, substantiated by a character, has to my knowledge been given it, in any botanical work, except the name of *Symplocarpus*, a term lately adopted by some American botanists on the alleged authority of Mr. Salisbury. As this name by its etymology implies a resemblance of the fruit to *Symplocos*, a genus with which the plant has not the least affinity; it appears to me inadmissible. Although I am averse to multiply the confusion

of synonyms, with which our science is already too much burdened, yet in the present instance an appropriate name, which should not be at variance with the character of the plant, appeared to be required. With the advice of the venerable Dr. Cutler, I have translated, as nearly as possible, the common English appellation for the plant. The name *Ictodes* from *ἴκτις, viverra*, and *οζω, oleo;* is sufficiently expressive of the property from which its common name is derived.

The genus Ictodes has for its character a *hooded spathe, spadix covered with perfect flowers, calyx with four segments, petals none, style pyramidal, seeds immersed in the spadix.* Only the present plant can be assigned to this genus. It belongs to *Tetrandria, monogynia;* and is found among the *Piperitæ* of Linnæus and *Aroideæ* of Jussieu. The root is large and abrupt, with numerous, crowded, fleshy fibres. The spathe which emerges from the ground some time before the leaves, is ovate, swelling, various in width, cucullate, spotted and sometimes nearly covered with dull brownish purple, the top acuminate and incurved, the edges infolded, auriculate at base, and at length coalescing. Within this is the oval spadix, on a short peduncle, covered with perfect tetrandrous flowers, and of the same colour with

JACOB BIGELOW *American Medical Botany* BOSTON, 1817-1820 CARY GRAPHIC ARTS COLLECTION, ROCHESTER INSTITUTE OF TECHNOLOGY Octavo